

## Disclaimer

- This product is a pharmaceutical product approved under a Japanese law, the Law for Ensuring the Quality, Efficacy and Safety of Drugs and Medical Devices, with a view to its sale and use in Japan.
- Multilingual product information is a translation of the product labeling written in Japanese and provided for your information only. It does not warrant that its contents and the product itself conforms to laws and regulations in countries other than Japan.
- Multilingual product information is a tentative translation by Our Company, and may be modified or altered without notice.
- Our Company assumes no responsibility for any occurred problem attributable to the contents of the multilingual product information.

# Sante<sup>®</sup> Medical Antibacterial

## Precautions for use

 **Things to avoid**  
(Not observing the following precautions may aggravate present symptoms or cause adverse reactions.)  
Do not use continuously for a long period of time.

## Consultation

1. The following people should consult with their doctor, pharmacist, or registered pharmaceutical distributor before using this product.
  - (1) People receiving treatment from a doctor
  - (2) People who have experienced an allergic reaction to medicine or other substances
  - (3) People who have the following symptom  
Severe pain in the eyes
2. If you experience any of the following symptoms after using this product, it may be an adverse reaction. In such a case, please stop using the product immediately, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.

| Area affected | Symptom                                     |
|---------------|---|
| Skin          | Rash/redness, itchiness                     |
| Eye(s)        | Redness, itchiness, swelling, stinging pain |

3. If symptoms do not improve within 3 to 4 days of use, please stop using the product, take this information form to your doctor, pharmacist, or registered pharmaceutical distributor and seek consultation.

## Dosage / Administration

Instill 1 drop at a time, 3 to 5 times daily.

### • Please observe the following precautions.

- (1) If administered to a child, the product must be administered under the direction and observation of the child's guardian.
- (2) Do not let the tip of the bottle touch your eyes, eyelids or eyelashes.
- (3) Do not use the product while wearing soft contact lenses.
- (4) Use only as an eye drop.




### Single-use type

Be sure to dispose of open vials immediately after use, even if there is some solution remaining.

### [Opening the bottle]

Slowly twist the cap to open it, without pulling on it forcefully.



## Indication

Sty, conjunctivitis (pink eye), blepharitis (sore eyelid), itchy eye

**Precautions regarding sty:** Be careful not to rub your eyelids or aggravate your eyes in other ways. (Some people may be susceptible to sty due to their constitution, but the condition can also easily form when the body is weakened due to overwork or lack of sleep. Please avoid fatigue and be sure to get sufficient sleep.)

## Ingredients and quantity

| Ingredients                            | Quantity | Action  |
|--|----------|---|
| Sulfamethoxazole (antibacterial agent) | 4.0%     | A long-acting sulfa drug that has an antibacterial action against the bacteria that cause conjunctivitis and sty. |
| Dipotassium glycyrrhizinate            | 0.25%    | Suppresses inflammation of the affected area.   |
| Taurine                                | 1.0%     | Activates cell metabolism of eye tissue.  |
| Vitamin B6                             | 0.1%     | Activates cell metabolism of eye tissue.  |

This product contains the following excipients: edetate sodium hydrate, hypromellose, boric acid, and pH-adjusting agent.

### [Precautions regarding ingredients and quantity]

You may experience either a bitter or a sweet taste in your mouth after putting the drops in your eyes.

This is due to the main active ingredients, sulfamethoxazole and dipotassium glycyrrhizinate, passing into your mouth through the lacrimal ducts. It is not due to degradation of quality or any other abnormalities of the product.

## Precautions regarding storage and handling

- (1) Store in a cool place away from direct sunlight.  
To preserve the quality of the product, do not leave it in a hot place such as inside a car or near a heater. If the product is left in a hot place, the vials may become warped causing the solution to leak and degrading its quality. Please do not use the product in such a case.
- (2) Keep out of reach of children.
- (3) Do not transfer the solution to another receptacle.  
(Doing so may lead to misuse of the product or may affect its quality.)
- (4) Do not share the product with another person.
- (5) To prevent light from altering the quality of the solution, it is contained in colored plastic vials sealed in a colored plastic bag. If storing the product, keep it in a place where it will not be exposed to light, such as by returning it to its box.
- (6) Placing the product into the back pocket of your trousers or other such places may cause the cap to open. Please exercise caution.
- (7) Do not use the product if it has exceeded its use-by date. Even before the use-by date, please use the product as soon as possible after opening the sealed bag.

